

PRACTICE EXAM 1 SOLUTIONS

Problem 1. For any set A , the empty set is an element of the power set of A .

Proof. This is true. The empty set is a subset of A , hence it is an element of the power set of A . □

Problem 2. For any sets A and B , we have $A - B \subseteq A$.

Proof. This is true. If $x \in A - B$ then $x \in A$ (and not in B). □

Problem 3. Let I be the set of natural numbers, and for each $i \in I$ let A_i be the closed interval in the real numbers $[1/i, i^2 + 1]$. Then

$$\bigcap_{i \in I} A_i = [1, 2].$$

Proof. This is true. The intervals are growing bigger as i increases, so their intersection is just $A_1 = [1, 2]$. □

Problem 4. Let $A = \{1, 2, 3\}$. Then A is a subset of the power set of A .

Proof. This is false. No element of A is a set, so they cannot belong to the power-set. □

Problem 5. If $a \equiv 3 \pmod{5}$, then $a^2 \equiv 4 \pmod{5}$.

Proof. This is true. Squaring both sides, we have $a^2 \equiv 3^2 = 9 \equiv 4 \pmod{5}$ since $5 \mid (9 - 4)$. □

Problem 6. Let A , B , and C be sets. Then $A - (B \cup C) = (A - B) \cap (A - C)$.

Proof. This is true. You can use Venn diagrams to see the equality. □

Problem 7. The converse of the statement “If x is even, then $x + 1$ is odd,” is the statement “If $x + 1$ is even, then x is odd.”

Proof. This is false. The given statement is the contrapositive, not the converse. □

Problem 8. The negation of the statement “There exists $x \in \mathbb{R}$, $x^2 - 1 < 0$,” is the statement “For all $x \in \mathbb{R}$, $x^2 - 1 < 0$.”

Proof. This is false. It should read “For all $x \in \mathbb{R}$, $x^2 - 1 \geq 0$.” □

Problem 9. The statement $P \wedge (\sim P)$ is a tautology.

Proof. This is false. You can see this using truth tables; this is a contradiction! □

Problem 10. Let A and B be sets. If A has seven elements, $A \cup B$ has ten elements, and $A - B$ has two elements, then B must contain eight elements.

Proof. This is true. Venn diagrams might help show you how many elements are in each set. □

Problem 11. For the following proof, determine which of the statements given below is being proved.

Proof. Assume a and b are odd integers. Then $a = 2k + 1$ and $b = 2\ell + 1$ for some $k, \ell \in \mathbb{Z}$. Then $ab^2 = (2k + 1)(2\ell + 1)^2 = 8k\ell^2 + 8k\ell + 2k + 4\ell^2 + 4\ell + 1 = 2(4k\ell^2 + 4k\ell + k + 2\ell^2 + 2\ell) + 1$. Since $4k\ell^2 + 4k\ell + k + 2\ell^2 + 2\ell \in \mathbb{Z}$, we see that ab^2 is odd. □

- a) If a or b is even, then ab^2 is even.
- b) If a and b are even, then ab^2 is even.
- c) If ab^2 is even, then a and b are even.
- d) If ab^2 is even, then a is even or b is even.
- e) None of the above.

Proof. The answer is (d). They are using the contrapositive. □

Problem 12. Let A be a set with 5 elements. Which of the following cannot exist:

- a) A subset of the power set of A with six elements.
- b) An element of the power set of A with six elements.
- c) An element of A containing six elements.
- d) Any of the above can exist, for suitable sets A .
- e) None of (a) through (c) can exist, no matter what A is.

Proof. The answer is (b) because elements of the power set are subsets of A , and subsets of A can have only elements of A . A subset of A can have at most 5 elements. □

Problem 13. Which of the following has a vacuous proof?

- a) Let $n \in \mathbb{Z}$. If $|n| < 1$ then $5n + 3$ is odd.
- b) Let $n \in \mathbb{Z}$. If $2n + 1$ is odd, then $n^2 + 1 > 0$.
- c) Let $x \in \mathbb{R}$. If $x^2 - 2x + 3 < 0$, then $2x + 3 < 5$.
- d) Let $x \in \mathbb{R}$. If $-x > 0$, then $x^2 + 3 > 3$.
- e) None of the above.

Proof. The answer is (c), because $x^2 - 2x + 3 = x^2 - 2x + 1 + 2 = (x - 1)^2 + 2 > 0$, so the premise is bogus. □

Problem 14. Which of the following statements has a trivial proof.

- a) Let $x \in \mathbb{N}$. If $x > 0$ then $x^2 > x$.
- b) Let $x \in \mathbb{N}$. If $x > 3$ then $2x$ is even.
- c) Let $x \in \mathbb{N}$. If $x < 2$ then $x^2 + 1$ is even.
- d) Let $x \in \mathbb{N}$. If $2x$ is even, then x is odd.

Proof. The answer is (b), since $2x$ is even, so the Q is true. □

Problem 15. Evaluate the following proof:

Theorem: Let $n \in \mathbb{Z}$. If $3n - 8$ is odd, then n is odd.

Proof. Let $n \in \mathbb{Z}$. Assume that n is odd. Then $n = 2k + 1$ for some integer k . Then

$$3n - 8 = 3(2k + 1) - 8 = 6k + 3 - 8 = 6k - 5 = 2(3k - 3) + 1.$$

Since $3k - 3 \in \mathbb{Z}$, we know that $3n - 8$ is odd. □

- a) The proof and the theorem are correct.
- b) The proof proves the converse of the given statement.
- c) The proof proves the contrapositive of the given statement.
- d) The proof contains arithmetic mistakes, which make it incorrect.
- e) None of the above.

Proof. The answer is (b). □

Problem 16. Let $A = \{\{1, 2\}, \{3, 4\}, \{5, 6\}\}$. The number of elements in the power set of A is

- a) 3 b) 4 c) 6 d) 8 e) 16 f) 64

Proof. The answer is (d). □

Problem 17. Let $x \in \mathbb{Z}$. The contrapositive of the open sentence “If x is even then $3x + 7$ is odd.” is the statement

- a) If x is odd then $3x + 7$ is even. b) If $3x + 7$ is odd then x is even.
 c) If $3x + 7$ is even then x is odd. d) If $3x + 7$ is even, then x is even.
 e) x is odd or $3x + 7$ is odd. f) x is odd or $3x + 7$ is even.

Proof. The answer is (c). □

Problem 18. Let x and y be integers. The negation of the statement “If xy is even then x is even or y is even” is

- a) If x is odd and y is odd, then xy is odd. b) If x is even or y is even, then xy is even.
 c) If xy is odd, then x is even and y is even. d) xy is even and x is odd and y is odd.
 e) xy is even and (x is odd or y is odd). f) xy is odd and (x is even or y is even).
 g) xy is odd and (x is odd and y is odd).

Proof. The answer is (d). Remember that the negation of an implication $P \Rightarrow Q$ is the statement $P \wedge \neg Q$. Also, the negation of an “or” is an “and”. □

Problem 19. If you wish to prove a statement of the form “If P then (Q or R)”, which of the following would **not** be a good way to begin.

- a) Assume P
 b) Assume $(\neg P) \wedge (Q \vee R)$
 c) Assume $(\neg Q) \wedge (\neg R)$.
 d) Assume $P \wedge (\neg Q) \wedge (\neg R)$.
 e) None of the above: all of these would be acceptable ways to begin.

Proof. The answer is (b). We never assume the negation of the premise when proving an implication. □

Problem 20. The following is a theorem proved in “Cohomology of number fields” (pg. 75) by J. Neukirch.

Theorem: Let G be a finite group, and let A, B be G -modules. If A is cohomologically trivial or B is divisible, then $\text{hom}(A, B)$ is cohomologically trivial.

Suppose that we know that G is a finite group, A and B are G -modules, and that $\text{hom}(A, B)$ is not cohomologically trivial. Which of the following must be true? (Think about the contrapositive.)

- a) A is cohomologically trivial and B is divisible.
 b) A is cohomologically trivial or B is divisible.
 c) A is not cohomologically trivial or B is divisible.
 d) A is not cohomologically trivial or B is not divisible.
 e) A is not cohomologically trivial and B is not divisible.

Proof. The answer is (e). □

Problem 21. Truth table.

Proof. Come see me if you need help on this one. □

Problem 22. Let $x, y \in \mathbb{Z}$. Prove that if $x^2 - xy$ is odd, then x is odd and y is even.

Proof. We prove the contrapositive. Assume x is even or y is odd.

Case 1: x is even. Then $x = 2k$ for some $k \in \mathbb{Z}$. Then $x^2 - xy = (2k)^2 - 2ky = 2(2k^2 - ky)$ is even since $2k^2 - ky \in \mathbb{Z}$.

Case 2: y is odd. We can also assume x is odd, else we are in case 1. Then $x = 2k + 1$ and $y = 2\ell + 1$ for some $k, \ell \in \mathbb{Z}$. Then $x^2 - xy = (2k + 1)^2 - (2k + 1)(2\ell + 1) = 4k^2 + 4k + 1 - 4k\ell - 2k - 2\ell - 1 = 2(k^2 + 2k - 2k\ell - k - \ell)$ is even since $k^2 + 2k - 2k\ell - k - \ell \in \mathbb{Z}$. \square

Problem 23. Prove the following statement. If x and y are rational, $x \neq 0$, and z is irrational, then $\frac{y+z}{x}$ is irrational.

Proof. Assume, by way of contradiction that $x, y \in \mathbb{Q}$, $x \neq 0$, z is irrational, and $\frac{y+z}{x} \in \mathbb{Q}$.

Since $x, \frac{y+z}{x} \in \mathbb{Q}$ their product $y + z = x \frac{y+z}{x}$ is rational.

Since $y + z, y \in \mathbb{Q}$, their difference $z = y + z - y$ is rational. This contradicts that fact that z is irrational. \square

Problem 24. Let $x, y \in \mathbb{R}$. Prove that if $x + y > 7$, then $x > 3$ or $y > 4$.

Proof. Let $x, y \in \mathbb{R}$. We work contrapositively. Assume $x \leq 3$ and $y \leq 4$. Thus $x + y \leq 3 + 4 = 7$. \square

Problem 25. Give examples of three sets A, B and C such that $A \in B$, $B \subseteq C$, and $A \not\subseteq C$.

Proof. Let $A = \{1\}$, $B = \{\{1\}\}$, and $C = B$. \square